




Rapid STAT®

Rapid STAT® is a **revolutionary point of collection test** for the detection of drugs of abuse in oral fluid, on surfaces or for pure substance testings.

Rapid STAT® combines the **convenience** of oral fluid collection, surface wipe testing or pure substance measurements with the **sensitivity, accuracy** and **precision** of a laboratory based test providing results in **about 6-8 minutes**.

NEW

One-Hand-Clip-System



AMP

Amphetamines 25 ng/ml

BZO

Benzodiazepines 25 ng/ml

COC

Cocaine 12 ng/ml

MTD

Methadone 15 ng/ml

MET♦MDMA

Methamphetamines 25ng/ml♦
Ecstasy 50 ng/ml

OPI

Opiates 25 ng/ml

PCP

Phencyclidine 2,5 ng/ml

THC

Marijuana 15 ng/ml

FLEXIBILITY

- 8 drugs available
- 2 to 6 drugs per test - varied configurations
- Can be applied as saliva-, wipe- or pure substance test

CONVENIENCE

- Easy to use due to the One-Hand-Clip-System
- All collections can be observed
- No restroom needed
- Specimen collection takes just 30 seconds
- Increased salivation due to innovative aroma field
- No shy bladder issues
- No specimen dilution or tampering issues
- **Results of saliva test ready in about 7-8 minutes**
- **Results of wipe- + pure substance testings ready in 6 min.**
- Rapid results allow for correct decisions

PERFORMANCE

- Each test strip has built-in procedural control
- Colloidal gold technology = superior sensitivity, accuracy & precision
- Tested for cross-reactivity against most common drugs and other substances
- Detects parent THC

SAMHSA states: "The only way to detect marijuana use is through the presence of the parent drug (THC) in the oral fluid....."

TRAINING/SUPPORT

- Onsite training
- Video Training on DVD
- Telephonic technical support



Minimum criteria for selecting oral fluid point of collection drug tests:

Rapid STAT® !

- 1. Device should test for the drugs recommended by SAMHSA**

The Substance Abuse Mental Health Services Administration (SAMHSA) has published proposed guidelines recommending that saliva based tests detect Amphetamines, Methamphetamines, MDMA, Cocaine, Opiates, PCP and THC.
- 2. Marijuana detection**

Many tests on the market are designed to detect THC metabolite (THC-COOH). Studies have shown that marijuana use will only be detected by testing for the residual parent drug. Therefore, detection of THC parent drug is required.
- 3. Sensitivity / cut-offs**

Drugs are present at much lower levels in saliva than in urine. Use a test that has the lowest cut-off levels to optimize identifying drug users.
- 4. Easy, rapid sample collection**

The collection of the saliva sample should be easy and should not take more than a few minutes.
- 6. Same samples used for confirmation**

The collection device should provide for a split sample, allowing the same sample to be sent for confirmation if necessary.
- 7. Training and support**

Oral fluid testing is relatively new, therefore training provided by the manufacturer should be comprehensive. Ongoing technical support is critical to maintaining an effective drug testing program. A manufacturer should provide both, training and support.